

### Cambridge Assessment International Education

Cambridge International Advanced Subsidiary and Advanced Level

| CANDIDATE<br>NAME |  |  |                     |  |  |
|-------------------|--|--|---------------------|--|--|
| CENTRE<br>NUMBER  |  |  | CANDIDATE<br>NUMBER |  |  |

CHEMISTRY 9701/31

Paper 3 Advanced Practical Skills 1

October/November 2019

2 hours

Candidates answer on the Question Paper.

Additional Materials: As listed in the Confidential Instructions

### **READ THESE INSTRUCTIONS FIRST**

Write your centre number, candidate number and name on all the work you hand in.

Give details of the practical session and laboratory where appropriate, in the boxes provided.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

Use of a Data Booklet is unnecessary.

Qualitative Analysis Notes are printed on pages 14 and 15.

A copy of the Periodic Table is printed on page 16.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [ ] at the end of each question or part question.

|   | Session    |
|---|------------|
|   |            |
|   |            |
|   | Laboratory |
|   |            |
| 1 |            |

| For Examiner's Use |  |  |
|--------------------|--|--|
| 1                  |  |  |
| 2                  |  |  |
| 3                  |  |  |
| Total              |  |  |

This document consists of 13 printed pages and 3 blank pages.



#### **Quantitative Analysis**

Read through the whole method before starting any practical work. Where appropriate, prepare a table for your results in the space provided.

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

1 In this experiment you will determine the concentration of a sample of hydrochloric acid. You will do this by measuring the volume of hydrogen produced when an excess of magnesium reacts with the acid.

$$Mg(s) + 2HCl(aq) \rightarrow MgCl_2(aq) + H_2(g)$$

**FA 1** is magnesium powder, Mg. **FA 2** is hydrochloric acid, HC *l*.

#### (a) Method

- Weigh the container with FA 1. Record the mass.
- Fill the tub with water to a depth of approximately 5 cm.
- Fill the 250 cm<sup>3</sup> measuring cylinder completely with water. Hold a piece of paper towel firmly over the top, invert the measuring cylinder and place it in the water in the tub.
- Remove the paper towel and clamp the inverted measuring cylinder so that the open end
  is just above the base of the tub.
- Use the 25 cm³ measuring cylinder to place 25.0 cm³ of **FA 2** into the reaction flask, labelled **X**.
- Check that the bung fits tightly in the neck of flask **X**, clamp flask **X**, and place the end of the delivery tube into the inverted 250 cm<sup>3</sup> measuring cylinder.
- Remove the bung from the neck of flask X. Tip all of FA 1 into flask X and replace the bung immediately. Remove the flask from the clamp and swirl to mix the contents.
- Swirl the flask occasionally until no more gas is evolved. Replace the flask in the clamp.
- Measure and record the final volume of gas in the measuring cylinder.
- Weigh and record the mass of the container with any residual solid.
- Calculate and record the mass of FA 1 used.

Keep FA 2 for use in Question 2.

| (b) | Cal        | culations                                                                                                                                  |
|-----|------------|--------------------------------------------------------------------------------------------------------------------------------------------|
|     | (i)        | Calculate the number of moles of hydrogen gas produced. (Assume 1 mol of gas occupies 24.0 dm³ at this temperature.)                       |
|     |            | moles of $H_2(g) = \dots mol$ [1]                                                                                                          |
|     | (ii)       | Calculate the concentration of hydrochloric acid in <b>FA 2</b> .                                                                          |
|     |            |                                                                                                                                            |
|     |            | concentration of HC $l$ in <b>FA 2</b> = mol dm <sup>-3</sup> [1]                                                                          |
| (   | (iii)      | In this experiment the magnesium powder was in excess.                                                                                     |
|     |            | Calculate the mass of magnesium powder needed for complete reaction with all the hydrochloric acid in $25.0\mathrm{cm^3}$ of <b>FA 2</b> . |
|     |            | mass of Mg =g<br>[1]                                                                                                                       |
| (c) |            | tudent suggested two modifications to the method in (a) to give a more accurate value for concentration.                                   |
|     | For        | each suggestion, state whether you agree with the student and explain your answer.                                                         |
|     | -          | ggestion 1: Use magnesium ribbon rather than powdered magnesium; keep the rest of the periment the same.                                   |
|     |            |                                                                                                                                            |
|     |            |                                                                                                                                            |
|     | Su(<br>san | ggestion 2: Use twice the mass of magnesium powder; keep the rest of the experiment the ne.                                                |
|     |            |                                                                                                                                            |
|     | ••••       | [2]                                                                                                                                        |

| Buy IGCSE, C | O / A Level Books, Past Papers & Revision Notes Online at Discounted Prices across Pakistan Visit: www.TeachifyMe.com / Shop |
|--------------|------------------------------------------------------------------------------------------------------------------------------|
| (d)          | Another student carried out the experiment in (a) but used less magnesium than that calculated in (b)(iii).                  |
|              | State and explain the effect this would have on the calculated concentration of hydrochloric acid in <b>FA 2</b> .           |
|              |                                                                                                                              |
|              | [1]                                                                                                                          |
|              | [Total: 8]                                                                                                                   |

5

2 In this experiment you will determine the concentration of **FA 2** by titration using aqueous sodium hydroxide.

$$HCl(aq) + NaOH(aq) \rightarrow NaCl(aq) + H2O(I)$$

**FA 2** is hydrochloric acid, HC*l*. **FA 3** is 0.100 mol dm<sup>-3</sup> sodium hydroxide, NaOH. methyl orange indicator

### (a) Method

#### Dilution of FA 2

- Fill the burette with **FA 2**.
- Run between 40.00 and 45.00 cm<sup>3</sup> from the burette into the 250 cm<sup>3</sup> volumetric flask.
- Record the volume used.
- Make the solution up to the 250 cm<sup>3</sup> mark by adding distilled water.
- Shake the flask thoroughly to ensure mixing.
- Label this solution of hydrochloric acid FA 4.

volume of **FA 2** used = ..... cm<sup>3</sup>

#### **Titration**

- Rinse the burette with distilled water and then with a little FA 4.
- Fill the burette with FA 4.
- Pipette 25.0 cm<sup>3</sup> of **FA 3** into a conical flask.
- Add several drops of methyl orange indicator.
- Perform a rough titration and record your burette readings.

The rough titre is ...... cm<sup>3</sup>.

- Carry out as many accurate titrations as you think necessary to obtain consistent results.
- Make sure any recorded results show the precision of your practical work.
- Record in a suitable form all of your burette readings and the volume of FA 4 added in each accurate titration.

| I    |  |
|------|--|
| II   |  |
| III  |  |
| IV   |  |
| V    |  |
| VI   |  |
| VII  |  |
| VIII |  |

[8]

|                     | your accurate titration results, obtain a value for the volume of <b>FA 4</b> to be used in your lations. Show clearly how you obtained this value. |
|---------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------|
|                     | 25.0 cm <sup>3</sup> of <b>FA 3</b> required cm <sup>3</sup> of <b>FA 4</b> . [1]                                                                   |
| (c) Calcı           | ulations                                                                                                                                            |
| (i) (               | Give your answers to (ii), (iii) and (iv) to the appropriate number of significant figures. [1]                                                     |
| (ii) C              | Calculate the number of moles of hydrochloric acid that reacted with 25.0 cm <sup>3</sup> of <b>FA 3</b> .                                          |
|                     | moles of HC <i>l</i> = mol [1]                                                                                                                      |
| (iii) (             | Calculate the concentration of hydrochloric acid in <b>FA 4</b> .                                                                                   |
|                     | concentration of HC $l$ in <b>FA 4</b> = mol dm <sup>-3</sup> [1]                                                                                   |
| (iv) (              | Calculate the concentration of hydrochloric acid in FA 2.                                                                                           |
|                     | concentration of HC $l$ in <b>FA 2</b> = mol dm <sup>-3</sup> [1]                                                                                   |
| (d) Calcu<br>flask. | ulate the maximum percentage error in the volume of FA 2 you added to the volumetric                                                                |
|                     | maximum percentage error = % [1]                                                                                                                    |

| (e) | In <b>Question 1</b> and <b>Question 2</b> you have determined the concentration of <b>FA 2</b> by two different methods. Each method used has possible sources of error, for example in <b>Question 1</b> the largest source of error is escape of gas. |
|-----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|     | Apart from this error, state and explain a source of error for each method.                                                                                                                                                                              |
|     | Question 1                                                                                                                                                                                                                                               |
|     |                                                                                                                                                                                                                                                          |
|     | Question 2                                                                                                                                                                                                                                               |
|     |                                                                                                                                                                                                                                                          |
|     | [2]                                                                                                                                                                                                                                                      |

[Total: 16]

Buy IGCSE, O / A Level Books, Past Papers & Revision Notes Online at Discounted Prices across Pakistan Visit: www.TeachifyMe.com / Shop \_ Call / WhatsApp: (0331-9977798)

### **Qualitative Analysis**

Where reagents are selected for use in a test, the **name** or **correct formula** of the element or compound must be given.

At each stage of any test you are to record details of the following:

colour changes seen;

3

- the formation of any precipitate and its solubility in an excess of the reagent added;
- the formation of any gas and its identification by a suitable test.

You should indicate clearly at what stage in a test a change occurs.

If any solution is warmed, a **boiling tube** must be used.

Rinse and reuse test-tubes and boiling tubes where possible.

No additional tests for ions present should be attempted.

| Qua  | alitative Analysis Notes.                                                                                                                                                                                                                            |
|------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| (i)  | Place a small spatula measure of <b>FA 5</b> in a hard-glass test-tube and heat <b>gently</b> . <b>Do not inhale the fumes.</b> Record <b>all</b> your observations.                                                                                 |
|      |                                                                                                                                                                                                                                                      |
|      |                                                                                                                                                                                                                                                      |
|      |                                                                                                                                                                                                                                                      |
|      | [2]                                                                                                                                                                                                                                                  |
| (ii) | Pour a 4 cm depth of distilled water into a boiling tube. Add the remaining <b>FA 5</b> and stir carefully until the solid has dissolved. This solution is <b>FA 6</b> .  Carry out the following tests on <b>FA 6</b> and record your observations. |

(a) FA 5 is a salt that contains two different cations and a single anion from those listed in the

| test                                                                  | observations |
|-----------------------------------------------------------------------|--------------|
| To a 1 cm depth in a test-tube, add aqueous ammonia.                  |              |
| To a 1 cm depth in a boiling tube, add aqueous sodium hydroxide, then |              |
| warm the mixture.                                                     |              |

| Buy IGCSE, O / A Level Books, Past Papers & Revision | Notes Online at Discounted Prices across Pakistan |
|------------------------------------------------------|---------------------------------------------------|
| Visit: www.TeachifyMe.com / Shop                     | Call / WhatsApp: (0331-9977798)                   |
| 9                                                    | , , ,                                             |

| test                                                                                        | observations |
|---------------------------------------------------------------------------------------------|--------------|
| To a 1 cm depth in a test-tube, add aqueous barium nitrate or aqueous barium chloride, then |              |
| add dilute hydrochloric acid or dilute nitric acid.                                         |              |

|    | Sanam Smeriae, alen                                                                                                                                 |                                                               |
|----|-----------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------|
|    | add dilute hydrochloric acid or dilute nitric acid.                                                                                                 |                                                               |
|    |                                                                                                                                                     | [4]                                                           |
| (i | ii) Identify the three ions in FA 5.                                                                                                                |                                                               |
|    | FA 5 contains, ,                                                                                                                                    | and                                                           |
|    | A student carried out Qualitative Analysis contained the ions $K^+$ , $Cr^{3+}$ and $SO_4^{2-}$ . The relative formula mass of <b>FA 7</b> is 499.3 | tests on a hydrated salt, <b>FA 7</b> , and concluded that it |
|    | Determine the formula of <b>FA 7</b> .                                                                                                              |                                                               |
|    |                                                                                                                                                     |                                                               |
|    |                                                                                                                                                     |                                                               |
|    |                                                                                                                                                     |                                                               |
|    | The formula of <b>FA 7</b> is                                                                                                                       | [2]                                                           |

(b)

Question 3 continues on page 10.

(c) FA 8 is a solution containing a single cation and a single anion, both of which are listed in the Qualitative Analysis Notes.

| ( | i) | Carry | out the | following | tests | and re | ecord | vour | observations |
|---|----|-------|---------|-----------|-------|--------|-------|------|--------------|
|   |    |       |         |           |       |        |       |      |              |

| test                                                                                                | observations |
|-----------------------------------------------------------------------------------------------------|--------------|
| To a 1 cm depth in a test-tube, add a few drops of aqueous acidified potassium manganate(VII), then |              |
| add starch indicator.                                                                               |              |
| To a 1 cm depth in a test-tube, add aqueous sodium hydroxide.                                       |              |

| a     | du staren mulcator.                                             |                                                           |
|-------|-----------------------------------------------------------------|-----------------------------------------------------------|
|       | o a 1 cm depth in a test-tube, add queous sodium hydroxide.     |                                                           |
|       |                                                                 | [2]                                                       |
| (ii)  | Identify the two ions in FA 8.                                  |                                                           |
|       | FA 8 contains                                                   | and[1]                                                    |
| (iii) | Suggest an additional test you could ca                         | arry out to confirm the presence of the anion in FA 8.    |
|       | Carry out this test and record your res                         | ult.                                                      |
|       |                                                                 |                                                           |
|       |                                                                 |                                                           |
|       |                                                                 | [2]                                                       |
| (iv)  | Give the ionic equation for the reaction Include state symbols. | n you carried out using <b>FA 8</b> and sodium hydroxide. |
|       |                                                                 | [1]                                                       |
|       |                                                                 |                                                           |

[Total: 16]

1.1

# **BLANK PAGE**

12

**BLANK PAGE** 

BLANK PAGE

# **Qualitative Analysis Notes**

# 1 Reactions of aqueous cations

| ion                                  | reaction with                                                                |                                                                              |  |  |  |  |  |  |  |
|--------------------------------------|------------------------------------------------------------------------------|------------------------------------------------------------------------------|--|--|--|--|--|--|--|
| ion                                  | NaOH(aq)                                                                     | NH <sub>3</sub> (aq)                                                         |  |  |  |  |  |  |  |
| aluminium,<br>A <i>l</i> ³+(aq)      | white ppt. soluble in excess                                                 | white ppt. insoluble in excess                                               |  |  |  |  |  |  |  |
| ammonium,<br>NH₄⁺(aq)                | no ppt. ammonia produced on heating                                          | _                                                                            |  |  |  |  |  |  |  |
| barium,<br>Ba <sup>2+</sup> (aq)     | faint white ppt. is nearly always observed unless reagents are pure          | no ppt.                                                                      |  |  |  |  |  |  |  |
| calcium,<br>Ca²+(aq)                 | white ppt. with high [Ca <sup>2+</sup> (aq)]                                 | no ppt.                                                                      |  |  |  |  |  |  |  |
| chromium(III),<br>Cr³+(aq)           | grey-green ppt. soluble in excess                                            | grey-green ppt. insoluble in excess                                          |  |  |  |  |  |  |  |
| copper(II),<br>Cu <sup>2+</sup> (aq) | pale blue ppt. insoluble in excess                                           | blue ppt. soluble in excess giving dark blue solution                        |  |  |  |  |  |  |  |
| iron(II),<br>Fe²+(aq)                | green ppt. turning brown on contact with air insoluble in excess             | green ppt. turning brown on contact with air insoluble in excess             |  |  |  |  |  |  |  |
| iron(III),<br>Fe³+(aq)               | red-brown ppt. insoluble in excess                                           | red-brown ppt. insoluble in excess                                           |  |  |  |  |  |  |  |
| magnesium,<br>Mg²+(aq)               | white ppt. insoluble in excess                                               | white ppt. insoluble in excess                                               |  |  |  |  |  |  |  |
| manganese(II),<br>Mn²+(aq)           | off-white ppt. rapidly turning brown on contact with air insoluble in excess | off-white ppt. rapidly turning brown on contact with air insoluble in excess |  |  |  |  |  |  |  |
| zinc,<br>Zn²+(aq)                    | white ppt. soluble in excess                                                 | white ppt. soluble in excess                                                 |  |  |  |  |  |  |  |

# 2 Reactions of anions

| ion                                            | reaction                                                                               |
|------------------------------------------------|----------------------------------------------------------------------------------------|
| carbonate,<br>CO <sub>3</sub> <sup>2-</sup>    | CO <sub>2</sub> liberated by dilute acids                                              |
| chloride,<br>C <i>l</i> <sup>-</sup> (aq)      | gives white ppt. with Ag <sup>+</sup> (aq) (soluble in NH <sub>3</sub> (aq))           |
| bromide,<br>Br <sup>-</sup> (aq)               | gives cream ppt. with Ag <sup>+</sup> (aq) (partially soluble in NH <sub>3</sub> (aq)) |
| iodide,<br>I-(aq)                              | gives yellow ppt. with Ag <sup>+</sup> (aq) (insoluble in NH <sub>3</sub> (aq))        |
| nitrate,<br>NO <sub>3</sub> -(aq)              | NH <sub>3</sub> liberated on heating with OH <sup>-</sup> (aq) and A <i>l</i> foil     |
| nitrite,<br>NO <sub>2</sub> -(aq)              | NH <sub>3</sub> liberated on heating with OH <sup>-</sup> (aq) and A <i>l</i> foil     |
| sulfate,<br>SO <sub>4</sub> <sup>2-</sup> (aq) | gives white ppt. with Ba <sup>2+</sup> (aq) (insoluble in excess dilute strong acids)  |
| sulfite,<br>SO <sub>3</sub> <sup>2-</sup> (aq) | gives white ppt. with Ba <sup>2+</sup> (aq) (soluble in excess dilute strong acids)    |

# 3 Tests for gases

| gas                             | test and test result                                                            |
|---------------------------------|---------------------------------------------------------------------------------|
| ammonia, NH <sub>3</sub>        | turns damp red litmus paper blue                                                |
| carbon dioxide, CO <sub>2</sub> | gives a white ppt. with limewater (ppt. dissolves with excess CO <sub>2</sub> ) |
| chlorine, Cl <sub>2</sub>       | bleaches damp litmus paper                                                      |
| hydrogen, H <sub>2</sub>        | 'pops' with a lighted splint                                                    |
| oxygen, O <sub>2</sub>          | relights a glowing splint                                                       |

| - 4 |   |
|-----|---|
|     |   |
|     | • |

|                                |       | 18 | 2   | <u>ө</u> | muil<br>O:      | 0. | <u>е</u> | neon<br>20.2     |               |               | 9.6                          | 9   | <u>ب</u> | pton<br>3.8       | 4  | (e)               | xenon<br>131.3     | 9,    | Rn               | uop -              |        |                   |                    |     |                  |                   |   |                    |
|--------------------------------|-------|----|-----|----------|-----------------|----|----------|------------------|---------------|---------------|------------------------------|-----|----------|-------------------|----|-------------------|--------------------|-------|------------------|--------------------|--------|-------------------|--------------------|-----|------------------|-------------------|---|--------------------|
|                                |       |    |     | т<br>—   | hel<br>4        | _  |          |                  |               |               |                              |     |          |                   |    |                   |                    |       |                  |                    |        |                   |                    |     |                  |                   |   |                    |
|                                |       | 17 |     |          |                 | 6  | ш        | fluorine<br>19.0 | 17            | Cl            | chlorine<br>35.5             | 35  | Ŗ        | bromine<br>79.9   | 53 | Ι                 | iodine<br>126.9    | 85    | Αŧ               | astatine<br>-      |        |                   |                    |     |                  |                   |   |                    |
|                                |       | 16 |     |          |                 | 80 | 0        | oxygen<br>16.0   | 16            | S             | sulfur<br>32.1               | 34  | Se       | selenium<br>79.0  | 52 | <u>e</u>          | tellurium<br>127.6 | 84    | Ъо               | polonium<br>—      | 116    | ^                 | livermorium<br>–   |     |                  |                   |   |                    |
|                                |       | 15 |     |          |                 | 7  | z        | nitrogen<br>14.0 | 15            | ₾             | phosphorus<br>31.0           | 33  | As       | arsenic<br>74.9   | 51 | Sb                | antimony<br>121.8  | 83    | <u>B</u>         | bismuth<br>209.0   |        |                   |                    |     |                  |                   |   |                    |
|                                |       | 14 |     |          |                 | 9  | ပ        | carbon<br>12.0   | 14            | S             | silicon<br>28.1              | 32  | Ge       | germanium<br>72.6 | 50 | Sn                | tin<br>118.7       | 82    | Pb               | lead<br>207.2      | 114    | Fl                | flerovium          |     |                  |                   |   |                    |
|                                |       | 13 |     |          |                 | 2  | В        | boron<br>10.8    | 13            | Αl            | aluminium<br>27.0            | 31  | Ga       | gallium<br>69.7   | 49 | In                | indium<br>114.8    | 81    | 11               | thallium<br>204.4  |        |                   |                    |     |                  |                   |   |                    |
|                                |       |    |     |          |                 |    |          |                  | •             |               | 12                           | 30  | Zu       | zinc<br>65.4      | 48 | පි                | cadmium<br>112.4   | 80    | Ρ̈́              | mercury<br>200.6   |        | ပ်                | copernicium        |     |                  |                   |   |                    |
| ments                          | Group |    |     |          |                 |    |          |                  |               |               |                              |     |          | 7                 | 29 | Cn                | copper<br>63.5     | 47    | Ag               | silver<br>107.9    | 62     | Au                | gold<br>197.0      | 111 | Rg               | roentgenium –     |   |                    |
| The Periodic Table of Elements |       |    |     |          |                 |    |          |                  |               |               |                              |     |          |                   |    |                   | 10                 | 28    | Z                | nickel<br>58.7     | 46     | Pd                | palladium<br>106.4 | 78  | 귙                | platinum<br>195.1 |   | Ds                 |
| riodic Ta                      |       |    |     |          |                 |    |          |                  |               |               |                              |     | 6        | 27                | ပိ | cobalt<br>58.9    | 45                 | 돈     | rhodium<br>102.9 | 77                 | 'n     | iridium<br>192.2  | 109                | ¥   | meitnerium<br>-  |                   |   |                    |
| The Pe                         |       |    | - : | I        | hydrogen<br>1.0 |    |          |                  |               |               | 80                           | 26  | Fe       | iron<br>55.8      | 44 | Ru                | ruthenium<br>101.1 | 92    | Os               | osmium<br>190.2    | 108    | Hs                | hassium            |     |                  |                   |   |                    |
|                                |       |    |     |          |                 |    |          | ,                |               |               |                              |     | 7        | 22                | Mn | manganese<br>54.9 | 43                 | ည     | technetium<br>-  | 75                 | Re     | rhenium<br>186.2  | 107                | В   | bohrium          |                   |   |                    |
|                                |       |    |     |          |                 |    |          |                  | atomic number | pol           | ass                          |     |          | 9                 | 24 | ပ်                | chromium<br>52.0   | 42    | Мо               | molybdenum<br>95.9 | 74     | ≯                 | tungsten<br>183.8  | 106 | Sg               | seaborgium<br>-   |   |                    |
|                                |       |    |     |          |                 |    |          | Key              |               | atomic symbol | name<br>relative atomic mass |     |          | 2                 | 23 | >                 | vanadium<br>50.9   | 41    | qN               | niobium<br>92.9    | 73     | Б                 | tantalum<br>180.9  | 105 | Ор               | dubnium           |   |                    |
|                                |       |    |     |          |                 |    |          |                  |               |               |                              | ato | rela     |                   |    | 4                 | 22                 | j     | titanium<br>47.9 | 40                 | Zr     | zirconium<br>91.2 | 72                 | Ξ   | hafnium<br>178.5 | 104               | 꿒 | rutherfordium<br>- |
|                                |       |    |     |          |                 |    |          |                  |               |               | က                            | 21  | Sc       | scandium<br>45.0  | 39 | >                 | yttrium<br>88.9    | 57–71 | lanthanoids      |                    | 89-103 | actinoids         |                    |     |                  |                   |   |                    |
|                                |       | 2  |     |          |                 | 4  | Be       | beryllium<br>9.0 | 12            | Mg            | magnesium<br>24.3            | 20  | Ca       | calcium<br>40.1   | 38 | Š                 | strontium<br>87.6  | 99    | Ba               | barium<br>137.3    | 88     | Ra                | radium             |     |                  |                   |   |                    |
|                                |       | _  |     |          |                 | 3  | :-       | lithium<br>6.9   | 7             | Na            | sodium<br>23.0               | 19  | $\prec$  | potassium<br>39.1 | 37 | Rb                | rubidium<br>85.5   | 55    | Cs               | caesium<br>132.9   | 87     | ъ                 | francium           |     |                  |                   |   |                    |

| 71 | 'n | lutetium<br>175.0     | 103 | ۲      | lawrencium   | ı     |  |
|----|----|-----------------------|-----|--------|--------------|-------|--|
| 70 | Υp | ytterbium<br>173.1    | 102 | 9<br>N | nobelium     | ı     |  |
| 69 | TB | thulium<br>168.9      | 101 | Md     | mendelevium  | ı     |  |
| 89 | щ  | erbium<br>167.3       | 100 | Fm     | ferminm      | ı     |  |
| 29 | 웃  | holmium<br>164.9      | 66  | Es     | einsteinium  | ı     |  |
| 99 | ò  | dysprosium<br>162.5   | 86  | Ç      | californium  | ı     |  |
| 65 | Д  | terbium<br>158.9      | 26  | Ř      | berkelium    | ı     |  |
| 64 | В  | gadolinium<br>157.3   | 96  | Cm     | curium       | ı     |  |
| 63 | Ē  | europium<br>152.0     | 92  | Am     | americium    | ı     |  |
| 62 | Sm | samarium<br>150.4     | 94  | Pu     | plutonium    | ı     |  |
| 61 | Pm | promethium<br>-       | 93  | dN     | neptunium    | ı     |  |
| 09 | PZ | neodymium<br>144.4    | 92  | ⊃      | uranium      | 238.0 |  |
| 59 | Ą  | praseodymium<br>140.9 | 91  | Ра     | protactinium | 231.0 |  |
| 28 | Se | cerium<br>140.1       | 06  | T      | thorium      | 232.0 |  |
| 25 | Гa | lanthanum<br>138.9    | 88  | Ac     | actinium     | ı     |  |

lanthanoids actinoids

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.