



## Cambridge International AS & A Level

---

**CHEMISTRY**

**9701/34**

Paper 3 Paper 34 (Advanced Practical Skills 2)

**May/June 2020**

MARK SCHEME

Maximum Mark: 40

---

### Published

Students did not sit exam papers in the June 2020 series due to the Covid-19 global pandemic.

This mark scheme is published to support teachers and students and should be read together with the question paper. It shows the requirements of the exam. The answer column of the mark scheme shows the proposed basis on which Examiners would award marks for this exam. Where appropriate, this column also provides the most likely acceptable alternative responses expected from students. Examiners usually review the mark scheme after they have seen student responses and update the mark scheme if appropriate. In the June series, Examiners were unable to consider the acceptability of alternative responses, as there were no student responses to consider.

Mark schemes should usually be read together with the Principal Examiner Report for Teachers. However, because students did not sit exam papers, there is no Principal Examiner Report for Teachers for the June 2020 series.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the June 2020 series for most Cambridge IGCSE™ and Cambridge International A & AS Level components, and some Cambridge O Level components.

---

This document consists of **9** printed pages.

**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

**GENERIC MARKING PRINCIPLE 1:**

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

**GENERIC MARKING PRINCIPLE 2:**

Marks awarded are always **whole marks** (not half marks, or other fractions).

**GENERIC MARKING PRINCIPLE 3:**

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

**GENERIC MARKING PRINCIPLE 4:**

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

**GENERIC MARKING PRINCIPLE 5:**

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

**GENERIC MARKING PRINCIPLE 6:**

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

**Science-Specific Marking Principles**

- 1 Examiners should consider the context and scientific use of any keywords when awarding marks. Although keywords may be present, marks should not be awarded if the keywords are used incorrectly.
- 2 The examiner should not choose between contradictory statements given in the same question part, and credit should not be awarded for any correct statement that is contradicted within the same question part. Wrong science that is irrelevant to the question should be ignored.
- 3 Although spellings do not have to be correct, spellings of syllabus terms must allow for clear and unambiguous separation from other syllabus terms with which they may be confused (e.g. ethane / ethene, glucagon / glycogen, refraction / reflection).
- 4 The error carried forward (ecf) principle should be applied, where appropriate. If an incorrect answer is subsequently used in a scientifically correct way, the candidate should be awarded these subsequent marking points. Further guidance will be included in the mark scheme where necessary and any exceptions to this general principle will be noted.

5 'List rule' guidance (see examples below)

For questions that require *n* responses (e.g. State **two** reasons ...):

- The response should be read as continuous prose, even when numbered answer spaces are provided
- Any response marked *ignore* in the mark scheme should not count towards *n*
- Incorrect responses should not be awarded credit but will still count towards *n*
- Read the entire response to check for any responses that contradict those that would otherwise be credited. Credit should **not** be awarded for any responses that are contradicted within the rest of the response. Where two responses contradict one another, this should be treated as a single incorrect response
- Non-contradictory responses after the first *n* responses may be ignored even if they include incorrect science.

6 Calculation specific guidance

Correct answers to calculations should be given full credit even if there is no working or incorrect working, **unless** the question states 'show your working'.

For questions in which the number of significant figures required is not stated, credit should be awarded for correct answers when rounded by the examiner to the number of significant figures given in the mark scheme. This may not apply to measured values.

For answers given in standard form, (e.g.  $a \times 10^n$ ) in which the convention of restricting the value of the coefficient (*a*) to a value between 1 and 10 is not followed, credit may still be awarded if the answer can be converted to the answer given in the mark scheme.

Unless a separate mark is given for a unit, a missing or incorrect unit will normally mean that the final calculation mark is not awarded. Exceptions to this general principle will be noted in the mark scheme.

7 Guidance for chemical equations

Multiples / fractions of coefficients used in chemical equations are acceptable unless stated otherwise in the mark scheme.

State symbols given in an equation should be ignored unless asked for in the question or stated otherwise in the mark scheme.

Question	Answer	Marks
1(a)	I Initial and final burette readings and volume added recorded for rough titre and accurate titre details tabulated.	1
	II Initial and final burette readings recorded and vol of <b>FB 3</b> added recorded for each accurate titration. Headings and units correct for accurate titrations. Headings: initial / final (burette) reading / volume <b>or</b> reading / volume at start / finish <b>and</b> volume / <b>FB 3</b> added/used or titre <b>and</b> units (cm <sup>3</sup> ) or / cm <sup>3</sup> <b>or</b> in cm <sup>3</sup> <b>or</b> cm <sup>3</sup> by every entry	1
	III All accurate burette readings are recorded to the nearest 0.05 cm <sup>3</sup> . Do not award this mark if: 50(.00) is used as an initial burette reading; More than one final burette reading is 50(.00) Any burette reading is > 50(.00)	1
	IV The final accurate titre recorded is within 0.10 cm <sup>3</sup> of any other accurate titre.	1
1(a)	<b>V, VI and VII</b> Award <b>V, VI and VII</b> for $\delta \leq 0.30 \text{ cm}^3$ Award <b>V and VI</b> for $0.30 \text{ cm}^3 < \delta \leq 0.50 \text{ cm}^3$ Award <b>V</b> for $0.30 \text{ cm}^3 < \delta \leq 0.80 \text{ cm}^3$	3
1(b)	Calculation of mean Candidate must average two (or more) accurate titres where the <b>total</b> spread is $\leq 0.20 \text{ cm}^3$ . Working must be shown or ticks must be put next to the two (or more) accurate readings selected.	1
1(c)(i)	all answers to <b>(ii)</b> , <b>(iii)</b> , <b>(iv)</b> and <b>(v)</b> to 3–4 sf (minimum 4 answers)	1
1(c)(ii)	Correctly calculates moles NaOH = $\frac{\text{(b)} \times 0.100}{1000}$ to 3–4 sf <b>and</b> same answer for moles of HCl	1

Question	Answer	Marks
1(c)(iii)	Correctly calculates moles $\text{HCl} = \frac{25.0 \times 0.200}{1000} = 5.00 \times 10^{-3}$ <b>and</b> moles neutralised = moles $\text{HCl}$ above – moles $\text{HCl}$ in (i)	1
1(c)(iv)	$\text{Na}_2\text{CO}_3(\text{aq}) + 2\text{HCl}(\text{aq}) \rightarrow 2\text{NaCl}(\text{aq}) + \text{CO}_2(\text{g}) + \text{H}_2\text{O}(\text{l})$ <b>and</b> correctly uses moles $\text{Na}_2\text{CO}_3 = \frac{\text{moles in HCl in (ii)}}{2}$	1
1(c)(v)	ans (iii) $\times 100$  $37.5 / ((\text{iii}) \times 100)$ can be done in either order  $\text{moles} = \frac{37.5}{100 \times (\text{iii})} = 2 \text{ marks}$ any error loses 1 mark	1
1(c)(vi)	Correctly calculates $x ((\text{Mr} - 106) / 18)$ and gives to nearest integer allow ecf even if negative integer.	1
1(d)	Disadvantage: if make mistake all the solution is wrong  Advantage: quicker/ larger volume has less % error	1

Question	Answer	Marks
2(a)	I Table with headings and units for 3 weighings and 3 calculated masses	1
	II Mass <b>FB 4</b> used between 1.80 and 2.20 g masses of <b>FB 4</b> , residue and water lost correctly calculated.	1
	III Ratio within 1.95–2.15 of supervisor ratio	1
	III and IV Ratio within 2.00–2.10 of supervisor ratio	1
2(b)(i)	Correctly calculates moles water = $\frac{\text{mass water}}{18}$ to 2–4sf Penalise sf only once in this question.	1
2(b)(ii)	$\text{MSO}_4 \cdot 7\text{H}_2\text{O}(\text{s}) \rightarrow \text{MSO}_4(\text{s}) + 7\text{H}_2\text{O}(\text{g})$ and correctly uses moles anhydrous = (ii) / 7	1
2(b)(iii)	Correctly uses mass $\frac{\text{FB 4}}{\text{moles anhydrous}} = M_r$	1
2(b)(iv)	Shows $A_r \text{ M} + 222.1 = M_r \text{ FB 4}$ and calculates $A_r$ of <b>M</b>	1
	identifies <b>M</b> as Group 2 metal with nearest $A_r$	1
2(c)(i)	To avoid water being picked up (from air)	1
2(c)(ii)	$A_r$ higher <b>with explanation</b> Mass water lost less $\rightarrow$ moles water lost less $\rightarrow$ moles anhydrous more $\rightarrow M_r$ higher $\rightarrow A_r$ higher very little effect because very little water vapour absorbed if lid prevented loss in (i) then allow $A_r$ lower as ecf	1

Question	Answer	Marks						
<b>FB 5</b> is Zn <b>FB 6</b> is ZnSO <sub>4</sub> .7H <sub>2</sub> O <b>FB 7</b> is propanone								
3(a)(i)	(Yellow solution goes) to blue / green	<b>1</b>						
	then to lilac / violet / mauve	<b>1</b>						
3(a)(ii)	any two from; <ul style="list-style-type: none"> <li>• blue (ignore green if before blue)</li> <li>• green / yellow</li> <li>• KMnO<sub>4</sub> loses purple colour</li> </ul>	<b>1</b>						
3(a)(iii)	Redox	<b>1</b>						
3(a)(iv)	Fizz / bubbles / effervescence	<b>1</b>						
	Gas / H <sub>2</sub> (gas) burns with a pop / pops with a lighted splint	<b>1</b>						
3(b)(i)	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 30%;">test</th> <th>observation</th> </tr> </thead> <tbody> <tr> <td><b>FB 6</b> + NaOH</td> <td>white ppt* soluble in excess*</td> </tr> <tr> <td><b>FB 6</b> + NH<sub>3</sub></td> <td>white ppt* soluble in excess *</td> </tr> </tbody> </table>	test	observation	<b>FB 6</b> + NaOH	white ppt* soluble in excess*	<b>FB 6</b> + NH <sub>3</sub>	white ppt* soluble in excess *	<b>2</b>
	test	observation						
	<b>FB 6</b> + NaOH	white ppt* soluble in excess*						
<b>FB 6</b> + NH <sub>3</sub>	white ppt* soluble in excess *							
Award one mark for every two correct observations (*)								
3(b)(ii)	<b>FB 5</b> is Zn / zinc (metal)	<b>1</b>						
3(b)(iii)	Zn(s) + H <sub>2</sub> SO <sub>4</sub> (aq) → ZnSO <sub>4</sub> (aq) + H <sub>2</sub> (g) ecf for incorrect <b>FB 5</b> provided gas is produced.	<b>1</b>						



Question	Answer	Marks
3(c)(i)	Choice of reagent: Tollens' and warming / Fehling's and warming / Benedict's and warming / Sandell's and warming / acidified potassium manganate(VII)	1
	Positive test for ethanal – silver mirror / black (or dark grey) solid for Tollens' Red / brown (solid / ppt) for Fehling's / Benedict's / Sandell's Purple to colourless for acidified manganate(VII) negative result for propanone – no reaction	1
3(c)(ii)	Negative test result and <b>FB 7</b> is propanone	1